

National Aeronautics and
Space Administration



ARSET

Applied Remote Sensing Training

<http://arset.gsfc.nasa.gov>

 @NASAARSET

Aerosol Observations from Satellites: Brief Theory & Existing Products

Pawan Gupta & ARSET Team

**Satellite Remote Sensing of Air Quality: Data, Tools,
and Applications**

Tuesday, May 23, 2017 – Friday, May 26, 2017

Indian Institute of Tropical Meteorology, Pune, India


Objectives

1. Gain a basic understanding of aerosol optical depth
2. Gain knowledge of and ability to access available aerosol products from NASA sensors

Aerosol Optical Depth

- AOD: Aerosol **Optical** Depth
- AOT: Aerosol **Optical** Thickness
- These optical measurements of light extinction are used to represent aerosol amounts in the entire column of the atmosphere

Optical Depth



The optical depth expresses the quantity of light removed from a beam by **scattering** or **absorption** during its path through a **medium**.

optical depth τ as


$$I = I_0 e^{-m\tau}$$

$$m = \sec \theta_0$$

$$\tau = \tau_{Rayl} + \tau_{aer} + \tau_{gas}$$

AERONET

<http://aeronet.gsfc.nasa.gov/>



Serves as a validation tool for satellite air quality products

Satellites for Air Quality Data

- MODIS (Terra and Aqua)
 - AOD: columnar aerosol loading – can be used to get PM_{2.5} or PM₁₀
- MISR (Terra)
 - Columnar aerosol loading in different particle size bins
 - In some cases aerosol heights
- OMI (Aura)
 - Absorbing aerosols, total aerosols
 - Trace gases
- VIIRS (NPP)
 - Aerosol optical depth
 - Aerosol type

Instrument Capabilities for Air Quality

Sensor Measurement Resolution

MODIS	250 m – 1 km
MISR	275 m – 1.1 km
OMI	13 x 24 km
VIIRS	750 m


A satellite image from the Moderate Resolution Imaging Spectroradiometer (MODIS) showing the Arctic region. The image displays a vast expanse of sea ice in the central Arctic Ocean, surrounded by landmasses including Alaska, Canada, and parts of Europe and Asia. The land is covered in green vegetation, brown tundra, and snow-capped mountains. The sea ice is shown in various shades of white and light blue, indicating different ice types and thicknesses. A semi-transparent grey rectangle is overlaid on the image, containing the text 'MODIS' and a horizontal line.

MODIS

Moderate Resolution Imaging Spectroradiometer

MODIS

- Spatial Resolution
 - 250 m, 500 m, 1 km
- Platform
 - Terra & Aqua
- Temporal Resolution
 - 2000-present
 - Daily, 8-day, 16-day, monthly, quarterly, yearly
- Data Format
 - Hierarchical Data Format – Earth Observing System Format (HDF-EOS)




- Spectral Coverage
 - 36 bands (major bands include red, blue, IR, NIR, MIR)
 - Bands 1-2: 250 m
 - Bands 3-7: 500 m
 - Bands 8-36: 1,000 m


A satellite image of Earth showing a large, hazy, brownish-grey aerosol plume originating from the west coast of Africa and spreading across the Atlantic Ocean. The plume is semi-transparent, allowing some land features to be seen underneath. The surrounding areas show typical satellite imagery of land with green vegetation and brown/tan dry areas, and blue oceans with white cloud patterns.

Aerosol Retrieval

Aerosol Detection



Complex Image: Smoke & Clouds




MODIS (Terra) May 14, 2007


Radiance to Aerosol Products



MODIS (Terra) May 2, 2007



Aerosol Retrieval Algorithm



Aerosol retrieval algorithm is a complex inversion scheme where assumptions are made in simulating satellite observations with advance radiative transfer calculations to retrieve atmospheric aerosol properties

Sources: Remer et al., 2005, Levy et al., 2013

MODIS Products

MOD01 Level-1A Radiance Counts

**MOD02 Level-1B Calibrated Geolocated Radiances –
also Level 1B”subsampled” 5kmx5km pro**

MOD03 Geolocation Data Set

MOD04 Aerosol Product

MOD05 Total Precipitable Water

MOD06 Cloud Products

MOD07 Atmospheric Profiles

MOD08 Gridded Atmospheric Product (Level 3)

MOD09 Atmospherically-corrected Surface Reflectance

MOD10 Snow Cover

MOD11 Land Surface Temperature & Emissivity

MOD12 Land Cover/Land Cover Change

MOD13 Vegetation Indices

MOD14 Thermal Anomalies, Fires & Biomass Burning

MOD15 Leaf Area Index & FPAR

MOD16 Surface Resistance & Evapotranspiration

**MOD17 Vegetation Production, Net Primary
Productivity**

MOD18 *Normalized Water-leaving Radiance

MOD19 Pigment Concentration

MOD20 Chlorophyll Fluorescence

MOD21 *Chlorophyll_a Pigment Concentration

MOD22 Photosynthetically Active Radiation (PAR)

MOD23 Suspended-Solids, Conc, Ocean Water

MOD24 Organic Matter Concentration

MOD25 Coccolith Concentration

MOD26 *Ocean Water Attenuation Coefficient

MOD27 Ocean Primary Productivity

MOD28 *Sea Surface Temperature

MOD29 Sea Ice Cover

MOD32 Processing Framework & Match-up Database

MOD33 Gridded Snow Cover

MOD34 Gridded Vegetation Indices

MOD35 Cloud Mask

MOD36 Total Absorption Coefficient

***MOD37 Ocean Aerosol Optical Thickness**

MOD39 Clear Water Epsilon

MOD43 Albedo 16-day L3

MOD44 Vegetation Cover Conversion

MYD – MODIS Aqua

MOD – MODIS Terra

A Few More Things About MODIS Data...

- MOD: Terra product
- MYD: Aqua product
- All MODIS products come in HDF format
- In HDF format each file contains both data and metadata
- **Scientific Data Set (SDS):** each parameter within a MODIS HDF file is referred to as an SDS
 - SDS must be referenced precisely according to name when analyzing the data within your own computer code


Things That Change with Each Instrument

(So you need to learn them!)

- Calibration Accuracy
- Quality Assurance – quality of the data
- Data Formats
- Product Resolutions
- Creating Level 3 products from Level 2
 - temporally and spatially averaging
- Current data release and data history

MODIS Aerosol Products


Three Separate Algorithms




- The dark target and deep blue products are separate and when both are available, the user must select which to use
- In collection 6, there is a joint product that uses an automated procedure to select the appropriate product

MODIS Aerosol Products


Two Algorithms




Dark Target





Deep Blue




Deep_Dark_Combined



MODIS 10km vs. 3km Products



High Resolution Aerosol Product



Quality Assurance is Extremely Important

QA indicates confidence in the quality of the retrieval

Quality_Assurance_Ocean

- Scale is 0-3
- Recommended Ocean QA above 1, 2, 3
- Factors:
 - Number of pixels
 - Error fitting
 - How close to glint

Quality_Assurance_Land


- Scale is 0-3
- Recommended Land QA of 3
- Factors:
 - Number of pixels
 - Error fitting
 - Surface reflectance

Understanding a MODIS File Name

Level 2, 10 km, Aerosol Product

Product Name

- Terra: MOD04
- Aqua: MYD04



HDFLook, Panoply, IDL, Python, Fortran, MatLab, and more can be used to read the data

Understanding a MODIS File Name

Level 2, 3 km, Aerosol Product

Product Name

- Terra: MOD04
- Aqua: MYD04



HDFLook, Panoply, IDL, Python, Fortran, MatLab, and more can be used to read the data


MODIS Aerosol Parameters (SDS)

- Optical_Depth_Land_and_Ocean
 - Retrieved using Dark Target Algorithm
 - Only high quality data
 - Over land QA = 3
 - Over ocean QA = 1, 2, 3
 - 10 km and 3km
- Dark_Target_Deep_Blue_Optical_Depth_550_Combined
 - Deep Blue & Dark Target Algorithm Merged Product
 - 10km only
- Dark_Target_Deep_Blue_Optical_Depth_550_Combined_QA
 - Quality flag associated with DD product

Access to MODIS Aerosol Products

- NASA LAADSWeb
 - Searchable database, FTP access
 - <http://ladsweb.nascom.nasa.gov/>
- MODIS-Atmos Site
 - Complete RGB archive with Level 3 product imagery
 - <http://modis-atmos.gsfc.nasa.gov>
- Giovanni for Level 3 Datasets
 - Web tool for imagery visualization and analysis
 - http://disc.gsfc.nasa.gov/gesNews/giovanni_3_end_of_service?instance_id=MODIS_DAILY_L3
- Dark Target Algorithm Site
 - <http://darktarget.gsfc.nasa.gov>
- Deep Blue Algorithm Site
 - <http://deepblue.gsfc.nasa.gov/>

Application of MODIS Aerosol Product



Source: van Donkelaar et al., 2006, 2009


Access to MODIS Aerosol Products

- NASA LAADSWeb
 - Searchable database, FTP access
 - <http://ladsweb.nascom.nasa.gov/>
- MODIS-Atmos Site
 - Complete RGB archive with Level 3 product imagery
 - <http://modis-atmos.gsfc.nasa.gov/>
- Giovanni for Level 3 Datasets
 - Web tool for imagery visualization and analysis
 - http://disc.gsfc.nasa.gov/gesNews/giovanni_3_end_of_service?instance_id=MODIS_DAILY_L3
- Dark Target Algorithm Site
 - <http://darktarget.gsfc.nasa.gov/>
- Deep Blue Algorithm Site
 - <http://deepblue.gsfc.nasa.gov/>



OMI

Ozone Monitoring Instrument (OMI)



Instrument Characteristics

- Nadir solar backscatter spectrometer
- Spectral Range: 270-500 nm
 - Resolution ~1 nm
- Swath Width: 2,600 km
 - Global daily coverage with 13x24 km spatial resolution

- One of four sensors on the EOS-Aura platform
 - OMI, MLS, TES, HIRDLS
- An international project
 - Holland, USA, Finland
- Launched July 15, 2004



Retrieval Products

- Column Amounts
 - Ozone (O_3)
 - Nitrogen Dioxide (NO_2)
 - Sulfur Dioxide (SO_2)
 - Others
- Aerosols

Applications of the Aerosol Index

Aerosols over clouds, April 14, 2006

- Validation tool for transport models
- Separation of carbonaceous from sulfate aerosols
- Tracking of aerosol plumes above clouds and over ice and snow




Above: Transport around the globe of a high altitude smoke layer generated by the Dec 2006 Australian fires. Numbers indicate the day of the month.



MISR

MISR Instrument




Source: Brian E.
Rheingans, JPL

Four MISR images over Appalachian Mountains
Nadir, 45.6 deg, 60.0 deg, 70.5 deg forward viewing cameras


Angular observations (which are not available in MODIS) makes MISR capable of providing additional information on particle size, shape and aerosol height under specific cases

MISR Global Daily Coverage



Applications of MISR Data

Smoke signals from the July 2004 Alaska and Yukon Fires



A satellite image of Earth showing a large semi-transparent gray rectangular box covering the central portion of the frame. The box contains the text 'VIIRS' and a horizontal line. The background image shows various geographical features: a large body of water on the left, a mountainous region with snow-capped peaks at the top, and a green, forested area on the right. The text 'VIIRS' is in a bold, black, sans-serif font.

VIIRS

Visible Infrared Imaging Radiometer (VIIRS)


A multi-wavelength imager like MODIS with similar wavelength bands

	MODIS	VIIRS
Orbit Altitude	690 km	824 km
Equator Crossing Time	13:30 LT	13:30 LT
Granule Size	5 min	86 sec
Swath	2,330 km	3,000 km
Pixel Nadir	0.5 km	0.75 km
Pixel Edge	2 km	1.5 km

VIIRS & MODIS


VIIRS

Nov 24, 2011







MODIS (Aqua)

Nov 24, 2011



CALIPSO: Vertical Profiles



Source: Meloë Kacenelenbogen

Satellite Aerosol Products

	MODIS	MISR	OMI	VIIRS
Strengths	Coverage Resolution Calibration Accuracy	Calibration Accuracy Particle shape Aerosol height for thick layer or plume	Indication of absorbing or scattering particles	Coverage Resolution Calibration Smaller bow-tie effect
Weaknesses	Bright Surfaces* Ocean glint Non-spherical particles	Coverage	Resolution Cloud contamination	Bright Surfaces* Ocean glint
Main Products	AOD Ocean–5 wavelengths Land–3 wavelengths Fine Fraction (Ocean only)	AOD 4 wavelengths Spherical/Non-spherical ratio Particle Size (3 Bins)	AOD AAOD Aerosol Index	AOD Aerosol Type
Product Resolution (level 2 and at Nadir)	10 Km 3 Km	17.6 Km	13 X 24 Km	0.75 km 6 km
Product Levels	2	2	2	2
Global Level 3 Aggregates	Daily 8 Day 30 Day	Monthly 3 Month Annual	Daily Monthly	Daily Monthly

Available Satellites for Aerosol Monitoring


	Pros	Cons
MODIS	<ul style="list-style-type: none"> • High spatial resolution (0.25-1 km) • Fine vs. coarse • Twice daily near-global coverage 	<ul style="list-style-type: none"> • No data under cloudy conditions • No vertical information • Larger uncertainties over bright targets
MISR	<ul style="list-style-type: none"> • Size/shape information • Higher accuracy • Multi-angle view 	<ul style="list-style-type: none"> • Limited swath width (360km) • Limited vertical information • No daily observations for air quality
OMI	<ul style="list-style-type: none"> • Daily near-global coverage • Absorbing aerosols • Precursor measurements (sulfate, nox) • Available over bright targets 	<ul style="list-style-type: none"> • Lack of information on scattering aerosols • Coarse resolution to separate clouds • Larger uncertainties
POLDER	<ul style="list-style-type: none"> • Daily near-global coverage • Sensitive to small mode aerosols • Available over bright targets 	<ul style="list-style-type: none"> • No data under cloudy conditions • No vertical information • Larger uncertainties over bright targets
CALIPSO	<ul style="list-style-type: none"> • Vertical information available • Information on clouds 	<ul style="list-style-type: none"> • Narrow swath (almost point measurement) • Very limited global coverage • Larger uncertainties in retrieved data sets
VIIRS, HIMAWARI, GOCI, and many more		

Satellite Limitations

- **Optical measurements**
 - only available in day time
 - very limited in night time
- Only available under
 - cloud free conditions
 - Snow/Ice free conditions
- **Accuracy** - vary (AOD) – Depends on satellite/algorithm
 - Very good over dark vegetated surfaces
 - Moderate over urban surfaces
 - Moderate to low over bright surface
 - Complex topography (i.e. mountains) – can be problematic
 - More uncertain for complex mixture of aerosols
- **Chemical Composition** - Very limited capabilities, only at research level
- **Temporal Coverage**
 - Usually once a day
 - But can use multiple satellite to get 2-3 a day
 - Geostationary will provide more frequent observations
- **Spatial Resolution**
 - 10 km (good)
 - 3 km (moderate)
 - 1 km, 0.75 km etc.

References & Links

- ARSET air quality page
 - <http://arset.gsfc.nasa.gov/airquality>
- NASA air quality
 - <http://airquality.gsfc.nasa.gov>
- MODIS Atmos
 - <http://modis-atmos.gsfc.nasa.gov/>
- MISR data
 - https://eosweb.larc.nasa.gov/PRODOCS/misr/Quality_Summaries/L2_AS_Products.html
- OMI data
 - <http://disc.sci.gsfc.nasa.gov/Aura/data-holdings/OMI>
- IDEA:
 - <http://www.star.nesdis.noaa.gov/smcd/spb/aq/>
- Smog blog:
 - <http://alg.umbc.edu/usaq/>

A satellite image of the North Atlantic Ocean, showing the eastern coast of North America on the left and the western coast of Europe on the right. The ocean is a deep blue, with some white clouds visible. A semi-transparent grey rectangular box is overlaid on the center of the image, containing the text "Questions & Discussion".

Questions & Discussion
